



Ph.D. thesis topics 2026/2027

DSP Rybářství / Fishery and DSP Ochrana vodních ekosystémů / Protection of aquatic ecosystems

Výzkumný ústav rybářský a hydrobiologický / Research Institute of fish Culture and Hydrobiology

Vodňany..... 2

Supervisor: Vitaliy Kholodnyy, Ph.D...... 3

Functional diversity of ovarian fluid in freshwater fish fertilization: effects on sperm performance, fertilization success, and species specificity..... 3

Supervisor: Prof. Otomar Linhart..... 5

Integrative analysis of the effects of short-term sperm storage on sperm physiology, embryonic proteome, and ncRNA-mediated transgenerational inheritance in common carp (*Cyprinus carpio*) and zebrafish (*Danio rerio*) 5

Supervisor: Jan Štundl, Ph.D...... 8

“Adult stem cells” and regeneration: The role of neural crest-derived Schwann cell precursors during the bone regeneration 8

Ústav akvakultury a ochrany vod / Institute of Aquaculture and Protection of Waters České

Budějovice..... 10

Supervisor: Quang Hung Tran, Ph.D...... 11

Deep Learning-Based Analysis of Feed Palatability in Aquaculture Species 11

Supervisor: Radek Gebauer, Ph.D...... 14

Sustainable Recovery and Valorization of Pond Sediment in Circular Bioeconomy 14

Supervisor: Koushik Roy, Ph.D...... 16

Carbohydrates in the pond food web, their interactions with fish nutrition and water quality 16

Supervisor: Assoc. prof. Vlastimil Stejskal..... 18

Effects of alternative feed ingredients on flesh quality, organoleptic properties, physicochemical characteristics and shelf life of farmed fish 18

Hydrobiologický ústav BC AV ČR / Institute of Hydrobiology CAS České Budějovice..... 21

Supervisor: Milan Říha, Ph.D...... 22

Spatiotemporal dynamics of fish movement: The role of orientation, temperature, and hydrology in habitat selection 22



Výzkumný ústav rybářský a hydrobiologický / Research Institute of fish Culture and Hydrobiology Vodňany





Supervisor: Vitaliy Kholodnyy, Ph.D.

Contact

E-mail: vkholodnyy@frov.jcu.cz

Tel.: +420 389 034 607

Functional diversity of ovarian fluid in freshwater fish fertilization: effects on sperm performance, fertilization success, and species specificity

Funkční diverzita ovariální tekutiny při oplození sladkovodních ryb: vliv na vlastnosti spermií, úspěšnost oplození a druhovou specifitu

DSP: Rybářství / Fishery

Annotation

Successful fertilization in externally fertilizing freshwater fish depends on highly coordinated interactions between spermatozoa, eggs, and the surrounding reproductive environment (Kholodnyy et al, 2020). Ovarian fluid (OF), the biologically active medium surrounding fish eggs during spawning, has traditionally been considered mainly as a protective or supportive substance. Recent studies indicate that OF may actively influence sperm motility parameters, longevity, swimming behavior, fertilization success, sperm longevity, and gamete interactions (Alonzo et al, 2016; Johnson et al, 2020) in a species-specific manner. Despite increasing interest in these processes, the biological functions and diversity of OF roles among freshwater fish species remain insufficiently understood.

Freshwater fish species differ markedly in their reproductive strategies, spawning environments, and susceptibility to natural or artificial hybridization. These differences may be reflected in the biochemical composition and physiological functions of OF (Gasparini, Evans 2018). Understanding these mechanisms is important not only for basic reproductive biology, but also for aquaculture, conservation programs, artificial reproduction, and the management of endangered or economically important fish species.

The proposed Ph.D. project will investigate how OF modulates sperm performance and fertilization processes in selected freshwater fish species, with particular focus on interspecific differences and intraspecific variability among females. The study will combine reproductive physiology, sperm motility analyses, in vitro fertilization experiments, and comparative biological approaches. Biochemical and molecular characterization of OF will be incorporated to identify factors associated with specific functional effects.

The main hypothesis

- Ovarian fluid actively modulates sperm motility patterns, swimming performance, and fertilization success in freshwater fish.
- The functional effects of ovarian fluid differ among fish species and among females within species.
- Ovarian fluid preferentially supports conspecific sperm and may contribute to species-specific fertilization and reproductive isolation.
- Functional differences in ovarian fluid are associated with biological, ecological, and biochemical variability among freshwater fish species.

Aim(s) of the Ph.D. thesis

- To characterize the effects of ovarian fluid on sperm performance parameters in selected freshwater fish species.



- To determine the influence of ovarian fluid on fertilization success under different experimental conditions.
- To investigate species-specific interactions between ovarian fluid and spermatozoa in conspecific and heterospecific combinations.
- To compare functional variability of ovarian fluid among fish species with different reproductive strategies.
- To identify selected biochemical or molecular characteristics associated with functional effects of ovarian fluid.

Possible approaches to reach the aims / to verify the hypotheses

- Collection of ovarian fluid, eggs, and sperm from selected freshwater fish species under controlled conditions.
- Evaluation of sperm motility and swimming behavior using Computer Assisted Sperm Analysis (CASA) systems under different activation conditions including ovarian fluid treatments.
- Performing fertilization experiments using homologous and heterologous ovarian fluid combinations under controlled laboratory conditions.
- Analysis of sperm longevity, swimming trajectories, velocity parameters, and fertilization outcomes in relation to ovarian fluid exposure.
- Comparative analysis of ovarian fluid effects among species differing in reproductive ecology and hybridization potential.
- Application of selected biochemical and molecular analyses of ovarian fluid including protein concentration measurements, ion composition analyses, metabolomic and proteomic approaches, or fractionation methods.
- Statistical analyses using generalized linear models (GLM), mixed-effects models, ANOVA, multivariate analyses, and correlation analyses.

References

- Alonzo, S.H., Stiver, K.A., Marsh-Rollo, S.E., 2016. Ovarian fluid allows directional cryptic female choice despite external fertilization. *Nat Commun* 7:12452. <https://doi.org/10.1038/ncomms12452>
- Gasparini, C., Evans, J.P., 2018 Female control over multiple matings increases the opportunity for postcopulatory sexual selection. *Proc Biol Sci.* 285 (1888): 20181505. <https://doi.org/10.1098/rspb.2018.1505>
- Johnson, S.L., Borziak, K., Keffmann, T., Rosengrave, P., Dorus, S., Gemmell, N.J., 2020. Ovarian fluid proteome variation associates with sperm swimming speed in an externally fertilizing fish, *J Evol Biol*, 33 (12): 1783–1794, <https://doi.org/10.1111/jeb.13717>
- Kholodnyy, V., Gadêlha, H., Cosson, J. and Boryshpolets, S., 2020. How do freshwater fish sperm find the egg? The physicochemical factors guiding the gamete encounters of externally fertilizing freshwater fish. *Rev Aquacult*, 12: 1165-1192. <https://doi.org/10.1111/raq.12378>

Funding

Czech Science Foundation (GAČR) project: “Decoding the Role of Female Reproductive Fluids in Freshwater Fish Fertilization”



Supervisor: Prof. Otomar Linhart

Contact

E-mail: linhart@frov.jcu.cz

Tel.: +420 724 357 897

Integrative analysis of the effects of short-term sperm storage on sperm physiology, embryonic proteome, and ncRNA-mediated transgenerational inheritance in common carp (*Cyprinus carpio*) and zebrafish (*Danio rerio*)

Integrovaná analýza vlivu krátkodobého uchování spermií na fyziologii spermií, embryonální proteom a transgenerační dědičnost zprostředkovanou nekódujícími RNA u kapra obecného (*Cyprinus carpio*) a zebřičky pruhované (*Danio rerio*)

DSP: Rybářství / Fishery

Annotation

Common carp (*Cyprinus carpio* L.) is one of the most important aquaculture species worldwide, whereas zebrafish (*Danio rerio*) represents a well-established vertebrate model organism widely used in developmental biology and biomedical research. Sperm contamination with urine is a major factor negatively affecting sperm quality during short-term storage in both species. Storage without an appropriate extender rapidly reduces sperm motility and fertilization capacity, while optimized storage conditions significantly improve sperm quality, fertilization success, and embryo hatching rates (Cheng et al., 2021a, 2022, 2024; Hossain et al., 2026; Ma et al., 2026a; Zhang et al., 2023, 2024, 2025).

Recent studies have demonstrated that carp sperm can be preserved for up to 30 days under optimized storage conditions (Ma et al., unpublished data). However, prolonged sperm storage induces adaptive changes in sperm membrane functionality and is associated with reduced motility, decreased fertilization capacity, and alterations in DNA methylation patterns (Cheng et al., 2025). These findings indicate that short-term sperm storage affects not only sperm physiological characteristics, but also paternal epigenetic information transmitted to the embryo. However, the identified epigenetic alterations represent the average of a heterogeneous sperm population, without distinguishing live-viable sperm subpopulations from dead sperm subpopulations (Cheng et al., 2021b), making it difficult to identify the real molecular responsible for alterations in embryonic development among the heterogeneous molecular signatures of the whole sperm population. In addition, the mechanisms linking storage-induced sperm non-coding RNAs (ncRNAs) with alterations in embryonic proteome and potential transgenerational effects remain largely unknown (Ma et al., 2026b). A comprehensive understanding of these processes is essential for optimizing reproductive biotechnology protocols and ensuring the production of high-quality gametes for sustainable aquaculture and biomedical applications. Integrating advanced cellular, epigenetic, transcriptomic, and proteomic approaches may provide new insights into how storage-induced sperm alterations influence early embryonic development and offspring performance.

The main hypothesis

- Short-term sperm storage differentially alters sperm the physiological properties of sperm subpopulations, including motility, plasma membrane integrity, mitochondrial activity, oxidative status, and DNA integrity.



- Characterization of changes in sperm membrane functionality will enable the optimization of short-term sperm storage and activation conditions, thereby preserving sperm fertilization potential.
- Physiological changes induced by short-term sperm storage led to alterations in the proteome of early embryos and affect biological processes associated with embryonic development.
- Short-term sperm storage differentially alters the profile of ncRNAs in sperm subpopulation, particularly miRNAs, piRNAs, and tRNA-derived small RNAs. The altered RNAs in live-viable sperm subpopulation can mediate the paternal transmission of epigenetic information to the embryo.
- Alterations in live-viable sperm ncRNAs profiles regulate gene expression and the proteome of early embryos, thereby influencing early embryonic development.
- Epigenetic changes induced by short-term sperm storage persist in the offspring and may be transmitted to subsequent generations through DNA methylation and ncRNAs-mediated inheritance, thereby affecting molecular regulation, development, and offspring phenotype.

Aim(s) of the Ph.D. thesis

- Characterization of the effects of short-term sperm storage on sperm quality, including motility. Characterization the differences between live-viable and dead sperm subpopulations, including membrane integrity, mitochondrial activity, oxidative status, and DNA integrity.
- Optimization of sperm storage and activation conditions based on changes in sperm membrane functionality to preserve fertilization capacity.
- Identification of changes in the embryonic proteome and biological pathways affected by paternal factors after fertilization with stored sperm.
- Characterization of storage-induced changes in sperm ncRNAs profiles and identification of candidate molecules involved in paternal epigenetic inheritance.
- Evaluation of the relationship between live-viable sperm ncRNAs alterations, embryonic gene regulation, and proteomic changes.
- Assessment of potential transgenerational effects of short-term sperm storage through monitoring of molecular, physiological, and phenotypic traits in offspring.
- Integration of physiological and multi-omics data to elucidate mechanisms linking sperm storage, embryonic development, and epigenetic inheritance.

Possible approaches to reach the aims / to verify the hypotheses

- Sperm quality assessment – analysis of motility (CASA), live-viable and dead sperm subpopulation separation, membrane integrity, mitochondrial activity, oxidative status, and DNA integrity using fluorescence-based methods and flow cytometry.
- Sperm storage optimization – comparison of storage and activation conditions using fertilization assays.
- Proteomic and ncRNA analyses – quantitative analysis of embryonic proteome (LC-MS/MS) and sequencing of sperm nRNAs (miRNA, piRNA, and tRNA-derived small RNAs).
- Data integration – bioinformatic analysis linking live-viable sperm ncRNA alterations with embryonic gene expression and proteomic changes.
- Transgenerational assessment – evaluation of molecular and phenotypic changes in F1 and, where possible, F2 generations, including identification of biomarkers associated with sperm subpopulation distinguish, and also epigenetic inheritance.

References

Cheng, Y., Franěk, R., Rodina, M., Xin, M.M., Cosson, J., Linhart, O., 2021 a. Optimization of sperm storage



- and fertilization in zebrafish *Danio rerio* (Hamilton). *Animals*, 11, 1558. <https://doi.org/10.3390/ani11061558>
- Cheng, Y., Vechtova, P., Fussy, Z., Sterba, J., Linhartová, Z., Rodina, M., Tučková V., Gela D., Samarin, A.M., Lebeda, I., Xin, M.M., Zhang, S., Rahi, D., Linhart, O., 2021 b. Changes in phenotypes and DNA methylation of in vitro aging sperm in common carp. *International Journal of Molecular Sciences*, 22, 5925, <https://doi.org/10.3390/ijms22115925>.
- Cheng, Y., Zhang, S., Linhartová, Z., Linhart, O., 2022. Common carp (*Cyprinus carpio*) sperm reduction during short-term in vitro storage at 4 °C. *Animal Reproduction Science*, 243, 107017. <https://doi.org/10.1016/j.anireprosci.2022.107017>
- Cheng, Y., Zhang, S., Nayak, R., Vechtova, P., Schumacher, F., Linhartová, P., Gazo, J., Linhartová, Z., Waghmare, S.G., Kleuser, B., Dey, A., Rodinová, V., Rodina, M., Sterba, J., Alavi, S.M.H., Labbé, C., Linhart, O., 2024. Fertilization by short-term stored sperm alters DNA methylation patterns at single-base resolution in common carp embryos. *Reviews in Fish Biology and Fisheries*. <https://doi.org/10.1007/s11160-024-09866-y>
- Cheng, Y., Alavi, S.M.H., Zhang, S., Nayak, R., Waghmare, S.G., Rahi, D., Ma, Z., Shazada, N.E., Linhart, O., Linhartová, Z., 2025. Comprehensive multi-omics analysis uncovers potential risks of aged sperm on offspring development after short-term storage. *BMC Biology*, 23, 264. <https://doi.org/10.1186/s12915-025-02379-5>
- Hossain, S., Ma, Z., Shazada, N.E., Zhang, S., Konar, E.S.M., Boryshpolets, S., Siddique, M.A.M., Shirupa, S., Policar, T., Linhart, O., 2026. Short-term preservation techniques for zebrafish (*Danio rerio*) sperm. *Animal Reproduction Science*, 292, 108254. <https://doi.org/10.1016/j.anireprosci.2026.108254>
- Ma, Z., Zhang, S., Hossain, S., Mohammad, Y., Alavi, S.M.H., Linhartová, Z., Dzyuba, V., Shazada, N.E., Gela, D., Linhart, O., 2026 a. Management of common carp (*Cyprinus carpio* L.) milt storage for genetic breeding programs: The role of extender and temperature. *Aquaculture Reports*, 48, 103602. <https://doi.org/10.1016/j.aqrep.2026.103602>
- Ma, Z., Zhang, S., Alavi, S.M.H., Linhartová, P., Cheng, Y., Boryshpolets, S., Dzyuba, B., Shazada, N.E., Linhart, O., 2026 b. The sperm RNA payload in fishes: A vector for paternal inheritance and memory. *Aquaculture Reports*, 47, 103490. <https://doi.org/10.1016/j.aqrep.2026.103490>
- Zhang, S., Cheng, Y., Alavi, S.M.H., Shazada, N.E., Linhartová, Z., Rodinová, V., Linhart, O., 2023. Elevated temperature promotes spermatozoa motility kinetics and fertilizing ability following short-term storage: An implication for artificial reproduction of common carp (*Cyprinus carpio*) in a hatchery. *Aquaculture*, 565, 739126. <https://doi.org/10.1016/j.aquaculture.2022.739126>
- Zhang, S., Bondarenko, O., Shazada, N.E., Cheng, Y., Rahi, D., Rodina, M., Gazi, I., Linhartová, Z., Siddique, M.A.M., Alavi, S.M.H., Boryshpolets, S., Dzyuba, B., Linhart, O., 2024. Plasma membrane contributes to temperature-induced spermatozoa motility kinetics in short-term stored sperm of common carp (*Cyprinus carpio*). *Aquaculture*, 583, 740588. <https://doi.org/10.1016/j.aquaculture.2024.740588>
- Zhang, S., Waghmare, S.G., Alavi, S.M.H., Shazada, N.E., Cheng, Y., Muthu, K.E.S., Vechtova, P., Brionne, A., Nayak, R., Ma, Z., Labbé, C., Linhart, O., Linhartová, Z., 2025. High temperature induces motility of zebrafish sperm after short-term storage. *Aquaculture*, 595, 741623. <https://doi.org/10.1016/j.aquaculture.2024.741623>

Funding

Decoding the role of female reproductive fluids in freshwater fish fertilization, 26-20614S, GAČR



Supervisor: Jan Štundl, Ph.D.

Contact

E-mail: jstundl@caltech.edu

Tel.: +1 (626) 696-7167

“Adult stem cells” and regeneration: The role of neural crest-derived Schwann cell precursors during the bone regeneration

„Dospělé kmenové buňky“ a regenerace: Role prekursorů Schwannových buněk odvozených z neurální lišty během regenerace kosti

DSP: Rybářství / Fishery

Annotation

The evolution of vertebrates is closely linked to the emergence of the neural crest, a migratory and multipotent stem cell population that gives rise to many vertebrate-specific features, including the craniofacial skeleton, peripheral neurons and glia, and numerous other cell types. However, it remains unclear whether neural crest stem cells persist in postnatal tissues and, if they do, whether they contribute to the regeneration of neural crest-derived tissues.

Among the neural crest-derived cell types present in postnatal tissues, Schwann cells and their precursors (SCPs) retain developmental properties similar to those of their embryonic maternal cells, neural crest stem cells. These cells are associated with peripheral nerves and, because of the extensive branching of the peripheral nervous system, are widely distributed throughout the body. Their regenerative roles following peripheral nerve injury are well documented, as Schwann cells are key contributors to nerve repair. However, peripheral nerves are not the only tissues in which neural crest-derived cells play important regenerative roles. It has been demonstrated that they are important for the regeneration of heart tissue, skin or teeth.

The bodies of sturgeons are covered by dermal bony scutes, neural crest-derived structures that represent an evolutionary remnant of the ancestral dermoskeleton of early vertebrates. Because these scutes retain a clear neural crest origin and possess regenerative capacity, sturgeons provide a unique opportunity to investigate whether SCPs contribute to the regeneration of neural crest-derived dermal bone.

The main hypothesis

- Schwann cell precursors represent an essential source of stem cell-like cells for the regeneration of dermal bones.

Aim(s) of the Ph.D. thesis

- To contribute to our understanding of Schwann cell precursors role in the regeneration of sturgeon bony scutes but also in general aspects of regeneration.

Possible approaches to reach the aims / to verify the hypotheses

- Analyses of bulk RNA-seq and scRNA-seq datasets.
- Validation of candidate genes using hybridization chain reaction in situ hybridization, immunohistochemistry.



- Functional validation using CRISPR/Cas9 mutagenesis.

References

- Stundl J, Martik ML, Chen D, Raja DA, Franěk R, Pospisilova A, Pšenička M, Metscher BD, Braasch I, Haitina T, Cerny R, Ahlberg PE, Bronner ME. Ancient vertebrate dermal armor evolved from trunk neural crest. *Proc Natl Acad Sci U S A*. 2023 Jul 25;120(30):e2221120120. doi: 10.1073/pnas.2221120120. Epub 2023 Jul 17. PMID: 37459514; PMCID: PMC10372632.
- Stundl J, Desingu Rajan AR, Bronner ME. Neural crest gene regulatory networks as drivers of development, diversification and disease. *Nat Rev Mol Cell Biol*. 2026 Jun;27(6):467-485. doi: 10.1038/s41580-026-00949-1. Epub 2026 Mar 4. PMID: 41781486; PMCID: PMC13085950.
- Johnston AP, Yuzwa SA, Carr MJ, Mahmud N, Storer MA, Krause MP, Jones K, Paul S, Kaplan DR, Miller FD. Dedifferentiated Schwann Cell Precursors Secreting Paracrine Factors Are Required for Regeneration of the Mammalian Digit Tip. *Cell Stem Cell*. 2016 Oct 6;19(4):433-448. doi: 10.1016/j.stem.2016.06.002. Epub 2016 Jul 1. PMID: 27376984.
- Kastri ME, Faure L, Von Ahsen D, Boudierlique TG, Boström J, Solovieva T, Jackson C, Bronner M, Meijer D, Hadjab S, Lallemend F, Erickson A, Kaucka M, Dyachuk V, Perlmann T, Lahti L, Krivanek J, Brunet JF, Fried K, Adameyko I. Schwann cell precursors represent a neural crest-like state with biased multipotency. *EMBO J*. 2022 Sep 1;41(17):e108780. doi: 10.15252/embj.2021108780. Epub 2022 Jul 11. PMID: 35815410; PMCID: PMC9434083.

Funding

GAČR Franěk 26-21447S



Fakulta rybnářství
a ochrany vod
Faculty of Fisheries
and Protection
of Waters

Jihočeská univerzita
v Českých Budějovicích
University of South Bohemia
in České Budějovice
Czech Republic

Ústav akvakultury a ochrany vod / Institute of Aquaculture and Protection of Waters

České Budějovice





Supervisor: Quang Hung Tran, Ph.D.

Contact

E-mail: htranquang@frov.jcu.cz

Tel. +420 389 034 682

Deep Learning-Based Analysis of Feed Palatability in Aquaculture Species

Analýza Chutnosti Krmiva U Akvakulturních Druhů Založená Na Hlubokém Učení

DSP: Rybářství / Fishery

Annotation

Over the past decade, the aquafeed industry has undergone a substantial transition from fishmeal and fish oil derived from marine forage fish to terrestrial protein sources and, more recently, to alternative and novel protein ingredients (Cottrell et al., 2020; Froehlich et al., 2018; Liu et al., 2025). This shift has reduced pressure on marine resources while supporting the expansion of aquaculture to meet the increasing global demand for seafood driven by population growth (Hua et al., 2019; Panteli et al., 2025).

Although alternative ingredients have been shown to maintain the production performance of aquaculture species at certain inclusion levels (Cottrell et al., 2020; Mugwanya et al., 2023; Qian et al., 2024; Tran et al., 2022), the palatability of these materials remains a challenge for feed acceptance in some fish species. Feed ingredients of aquatic animal origin, such as fishmeal, are rich in chemical compounds including free amino acids, nucleotides, amines, and nucleosides, which are readily detected by the chemosensory systems of aquatic animals during food detection and ingestion. In contrast, alternative protein sources are generally lower levels of these compounds, limiting their capacity to fully replace fishmeal in aquaculture feeds (Dong et al., 2016; Soengas et al., 2025; Suresh et al., 2011; Terrey et al., 2021). Therefore, improving the inclusion of unconventional protein sources in aquafeeds while maintaining feed acceptance requires the selection of suitable ingredients for specific species, additional ingredient processing, or the incorporation of feed attractants (Alves et al., 2020; Dong et al., 2016; Hossain et al., 2024; Terrey et al., 2021).

Feed palatability in aquaculture species has been evaluated using various assessment methods, including manual observation in multi-chamber tanks (Suresh et al., 2011), Y-maze systems (Jingting et al., 2020), self-feeding approaches (Roy et al., 2020), and hydrophone-based monitoring (Araujo et al., 2026). More recently, deep-learning-based techniques have emerged as valuable tools for investigating feed palatability in aquatic animals (Janssens et al., 2026; Tran et al., 2026). These approaches enable the analysis of behavioural indicators such as swimming speed, pellet ingestion dynamics, movement heatmaps, and feeding responses. However, the palatability of feeds containing alternative proteins and attractants has so far been insufficiently investigated using precise and lightweight deep learning approaches in the context of unconventional feed ingredients.



The main hypothesis

- The palatability of unconventional feed ingredients varies considerably in aquatic species.
- The use of attractants may enhance the palatability of feed ingredients with low acceptability.
- Free amino acids, nucleotides, and amines are the primary drivers of feed palatability.

Aim(s) of the Ph.D. thesis

- To evaluate feed palatability through fish behavior while fed unconventional protein sources, using a deep learning technique.
- To assess different commercial feed attractants on fish behavior using a deep learning technique
- To evaluate whether a specific attractant enhances the palatability of a low-palatability feed ingredient.

Possible approaches to reach the aims / to verify the hypotheses

- Three experiments will be conducted.
 - Experiment 1: A feeding trial will evaluate three protein sources, insect meal, fungal meal, and bacterial meal, as replacements for fishmeal in comparison with a control diet.
 - Experiment 2: A second feeding trial will evaluate three feed attractants in comparison with a control diet without attractants.
 - Experiment 3: One of the low-palatability diets identified in Experiment 1 will be used as a reference diet, while the remaining diets will be supplemented with a promising candidate identified in Experiment 2.
- In these experiments, the tested animals will be maintained under both normal and stressful conditions, such as hypoxia and crowding. Tank designs will range from conventional tanks to Y-maze and multi-chamber systems. Feed intake and fish behavior will be analyzed using a deep learning approach as previously described (Tran et al., 2026). Diets and ingredients will be analyzed for nutritional composition and specific palatability-related compounds. Fish will also be evaluated for physiological responses, including neurotransmitters, taste receptors, and related parameters.

References

- Alves, D.R.S., de Oliveira, S.R., Luczinski, T.G., Boscolo, W.R., Bittencourt, F., Signor, A., Detsch, D.T., 2020. Attractability and palatability of liquid protein hydrolysates for Nile tilapia juveniles. *Aquaculture Research*. 51 (4), 1681-1688. <https://doi.org/10.1111/are.14514>.
- Araujo, A.N., Peixoto, S., Davis, D.A., 2026. Effect of protein intake on growth, protein retention, feed consumption and acoustic feeding behavior of Pacific white shrimp (*Litopenaeus vannamei*) in clear water RAS. *Aquaculture*. 614 743547. <https://doi.org/10.1016/j.aquaculture.2025.743547>.
- Cottrell, R.S., Blanchard, J.L., Halpern, B.S., Metian, M., Froehlich, H.E., 2020. Global adoption of novel aquaculture feeds could substantially reduce forage fish demand by 2030. *Nature Food*. 1 (5), 301-308. 10.1038/s43016-020-0078-x.
- Dong, C., He, G., Mai, K., Zhou, H., Xu, W., 2016. Palatability of water-soluble extracts of protein sources and replacement of fishmeal by a selected mixture of protein sources for juvenile turbot (*Scophthalmus maximus*). *Journal of Ocean University of China*. 15 (3), 561-567. 10.1007/s11802-016-2898-8.
- Froehlich, H.E., Jacobsen, N.S., Essington, T.E., Clavelle, T., Halpern, B.S., 2018. Avoiding the ecological



- limits of forage fish for fed aquaculture. *Nature Sustainability*. 1 (6), 298-303. 10.1038/s41893-018-0077-1.
- Hossain, M.S., Small, B.C., Kumar, V., Hardy, R., 2024. Utilization of functional feed additives to produce cost-effective, ecofriendly aquafeeds high in plant-based ingredients. *Reviews in Aquaculture*. 16 (1), 121-153. <https://doi.org/10.1111/raq.12824>.
- Hua, K., Cobcroft, J.M., Cole, A., Condon, K., Jerry, D.R., Mangott, A., Praeger, C., Vucko, M.J., Zeng, C., Zenger, K., Strugnell, J.M., 2019. The Future of Aquatic Protein: Implications for Protein Sources in Aquaculture Diets. *One Earth*. 1 (3), 316-329. 10.1016/j.oneear.2019.10.018.
- Janssens, T., Nguyen, Tram T., Stejskal, V., Aebischer, P., Tran, Hung Q., 2026. Deep learning-based analysis of feed preference in Pacific white shrimp (*Litopenaeus vannamei*) reared in Y-maze tanks.
- Jingting, Y., Danting, G., Chun, K., Min, J., Xueming, H., 2020. Effect of soybean antigenic protein on feed palatability of fishmeal replaced diets for obscure puffer (*Takifugu fasciatus*) and the alternation of diet preference by domestication. *Aquaculture Reports*. 17 100332. <https://doi.org/10.1016/j.aqrep.2020.100332>.
- Liu, Y., Jiang, Z., Cottrell, R.S., Tlustý, M.F., Fitzsimmons, K., Cao, L., 2025. Unstable supply and future shortages of wild forage fish heighten risks to global fed aquaculture production. *Nature Food*. 6 (11), 1068-1078. 10.1038/s43016-025-01254-4.
- Mugwanya, M., Dawood, M.A.O., Kimera, F., Sewilam, H., 2023. Replacement of fish meal with fermented plant proteins in the aquafeed industry: A systematic review and meta-analysis. *Reviews in Aquaculture*. 15 (1), 62-88. <https://doi.org/10.1111/raq.12701>.
- Panteli, N., Kousoulaki, K., Antonopoulou, E., Carter, C.G., Nengas, I., Henry, M., Karapanagiotidis, I.T., Mente, E., 2025. Which Novel Ingredient Should be Considered the “Holy Grail” for Sustainable Production of Finfish Aquafeeds? *Reviews in Aquaculture*. 17 (1), e12969. <https://doi.org/10.1111/raq.12969>.
- Qian, Y.-F., Limbu, S.M., Qiao, F., Luo, Y., Chen, L.-Q., Zhang, M.-L., Du, Z.-Y., 2024. Seeking the best alternatives: A systematic review and meta-analysis on replacing fishmeal with plant protein sources in carnivorous fish species. *Reviews in Aquaculture*. 16 (3), 1099-1126. <https://doi.org/10.1111/raq.12888>.
- Roy, J., Mercier, Y., Tonnet, L., Burel, C., Lanuque, A., Surget, A., Larroquet, L., Corraze, G., Terrier, F., Panserat, S., Skiba, S., 2020. Rainbow trout prefer diets rich in omega-3 long chain polyunsaturated fatty acids DHA and EPA. *Physiology & Behavior*. 213 112692. <https://doi.org/10.1016/j.physbeh.2019.112692>.
- Soengas, J.L., Comesaña, S., Blanco, A.M., Conde-Sieira, M., 2025. Feed Intake Regulation in Fish: Implications for Aquaculture. *Reviews in Fisheries Sc*

Funding

Collaboration with Feed ingredient and Aquafeed Producers and laboratory sources.

CENAKVA Research program

RP3 Sustainable aquaculture with a responsible water and nutrient management



Supervisor: Radek Gebauer, Ph.D.

Contact

E-mail: rgebauer@frov.jcu.cz

Tel. +420 389 034 647

Sustainable Recovery and Valorization of Pond Sediment in Circular Bioeconomy

Udržitelné využití a zhodnocení rybníčního sedimentu v podmínkách cirkulární bioekonomiky

DSP: Rybníkářství / Fishery

Annotation

In Central Europe, fishpond management is associated with the long-term accumulation of organic matter, nutrients, suspended solids, and various contaminants originating from fish production, agricultural runoff, and surrounding catchments. Regular sediment removal is necessary to maintain pond functionality, water quality, and ecological stability. However, dredged pond sediments are frequently classified as waste materials, creating economic and environmental burdens related to their disposal and management. At the same time, these sediments contain substantial amounts of organic carbon, nitrogen, phosphorus, and micronutrients, indicating significant potential for reuse within circular bioeconomy systems (Drozd et al., 2020).

Increasing environmental pressure, stricter European legislation, and the need to reduce dependency on mineral fertilizers highlight the urgent necessity to develop safe and sustainable strategies for sediment valorization. Current knowledge concerning the environmental risks, treatment efficiency, and long-term impacts of sediment application on soil and aquatic ecosystems remains fragmented and insufficiently standardized. Furthermore, there is limited understanding of how treatment technologies such as composting, anaerobic digestion, pyrolysis, hydrothermal carbonization, or biochar stabilization influence contaminant mobility, nutrient availability, and the overall environmental footprint of sediment utilization (Drozd et al., 2020; Marval et al. 2020)).

Within this framework, pond sediments could become an important source of bio-based fertilizers, soil amendments, substrates for ecological restoration, or raw materials for industrial applications (Drozd et al., 2020). However, successful implementation requires comprehensive evaluation of sediment quality, environmental safety, treatment technologies, economic feasibility, and stakeholder acceptance.

The main hypothesis

- Appropriately treated pond sediments can serve as a safe and effective secondary resource for nutrient recovery and agricultural applications within circular bioeconomy systems.
- Advanced sediment treatment technologies (e.g., composting, pyrolysis, hydrothermal carbonization, or biochar stabilization) significantly reduce the environmental risks associated with pond sediments, including contaminant mobility.
- The valorization of pond sediments through circular bioeconomy approaches provides measurable environmental and economic benefits compared to conventional sediment disposal, including reduced waste generation, improved nutrient recycling efficiency, and lower environmental footprint.



Aim(s) of the Ph.D. thesis

- To characterize the physicochemical, microbiological, and contaminant profile of pond sediments originating from different aquaculture management systems, with emphasis on nutrient content, heavy metals, and micropollutants.
- To evaluate and optimize sustainable treatment and valorization technologies for pond sediments, including biological, thermochemical, and stabilization approaches, in order to improve nutrient availability and reduce environmental and health risks.
- To assess the environmental, agronomic, and economic potential of treated pond sediments within circular bioeconomy applications using integrated sustainability indicators, including nutrient recovery efficiency, ecotoxicological safety, and life cycle assessment (LCA).

Possible approaches to reach the aims / to verify the hypotheses

- Collection of sediment samples from different aquaculture systems followed by physicochemical, microbiological, and contaminant analyses, including nutrients, organic matter, heavy metals, micropollutants.
- Testing of treatment and stabilization technologies: Evaluation and optimization of sustainable sediment treatment methods such as composting, vermicomposting, anaerobic digestion, pyrolysis, hydrothermal carbonization, and biochar-based stabilization to reduce environmental risks and improve nutrient recovery.
- Assessment of agronomic and environmental safety: Performance of germination, phytotoxicity, and plant growth experiments to evaluate nutrient availability, soil improvement potential, contaminant mobility, and possible ecological risks associated with sediment application.
- Sustainability and circular bioeconomy evaluation: Application of life cycle assessment (LCA), nutrient flow analysis, and economic assessment to compare conventional sediment disposal with circular valorization pathways and identify sustainable utilization strategies.

References

- Drozd, D., Malinska, K., Mazurkiewicz, J., Kacprzak, M., Mrowiec, M., Szczypiór, A., Postawa, P. and Stachowiak, T., 2020. Fish pond sediment from aquaculture production-Current practices and the potential for nutrient recovery: a Review. *International Agrophysics*, 34(1).
- Marval, Š., Hejduk, T. and Zajíček, A., 2020. An analysis of sediment quality from the perspective of land use in the catchment and pond management. *Soil and Sediment Contamination: An International Journal*, 29(4), pp.397-420.

Funding

NAZV PFAS3AKVA (Gebauer PI), Interreg Aquacycle (Malinovskyi PI, Gebauer responsible for Laboratory of Nutrition)

CENAKVA Research program

RP3 Sustainable aquaculture with a responsible water and nutrient management



Supervisor: Koushik Roy, Ph.D.

Contact

E-mail: kroy@frov.jcu.cz

Tel. +420 389 034 648

Carbohydrates in the pond food web, their interactions with fish nutrition and water quality

Sacharidy v potravní síti rybníka, jejich interakce s výživou ryb a kvalitou vody

DSP: Rybnářství / Fishery

Annotation

Carbohydrates constitute one of the major macronutrient fractions of natural food web components in pond aquaculture, including algae, zooplankton and benthic invertebrates. Their nutritional value depends not on total abundance but on carbohydrate typology, digestibility and fermentability. The composition, availability, digestibility and bioenergetic significance of aquatic carbohydrates in pond food web remain largely unexplored from a fish nutritional point of view, limiting the development of precision pond feeds. Natural food organisms frequently contain large proportions of recalcitrant carbohydrates, collectively termed non-starch polysaccharides (NSPs), including chitin, cellulose, hemicellulose, pectin, lignin and β -glucans, which are generally resistant to digestion by fish. For example, carbohydrates in zooplankton and benthic invertebrates are often dominated by chitin, whereas aged algal and microbial biomasses may contain substantial quantities of cellulose and hemicellulose. This creates a mismatch between carbohydrate abundance and digestible non-protein energy availability within pond food webs. Such mechanisms partly explain the long-standing success of traditional grain-fed European carp ponds, where supplementary cereals provide readily digestible carbohydrates and non-protein energy that was otherwise limited in natural food/ pond ecosystem. Under conditions where lipid-rich natural food resources do not dominate carp diets, digestible carbohydrates do represent an important source of non-protein energy. The pond-reared common carp (*Cyprinus carpio*) may break down complex NSP fractions through synergistic digestibility effect of pond ecosystem and extract energy. But beyond energy supply, the interplay of carbohydrates type may strongly influence excretion, gut fullness, lipid digestibility, protein sparing, feed intake regulation, grazing pressure on natural food and nutrient cycling within pond ecosystems.

The main hypothesis

- Carbohydrate types including their stoichiometries represent an overlooked determinant of fish growth, supplementary feed efficiency, natural food grazing pressure and water quality in pond aquaculture.

Aim(s) of the Ph.D. thesis

- To investigate the seasonal dynamics of carbohydrate fractions and non-protein energy supplied by carp gill-filterable plankton communities and zoobenthic organisms throughout the vegetative season in temperate European fishponds.



- To investigate the interactions between different non-starch polysaccharide (NSP) fractions and the digestion/ absorption of other nutrients in fish, with particular emphasis on highly demanding digestion process of lipids.
- To determine the optimal carbohydrate composition of supplementary feeds for enhancing zooplankton abundance, improving fish bioenergetic efficiency, and reducing eutrophication risk in pond ecosystems.
- To quantify the capacity of pond aquaculture systems to convert recalcitrant carbohydrate inputs into human-edible fish biomass without increasing eutrophication or harmful algal bloom risk, relative to traditional grain-feeding systems.

Possible approaches to reach the aims / to verify the hypotheses

- Fish digestibility and nutritional bioenergetics trial in Guelph-RAS system.
- Pond trial with plankton net, Eckman grab operation.
- Pond water mesocosms, aquaponic sludge digestors (bioreactors).
- FOSS FibreTec analyses (for NDF, ADF, ADL analyses).
- LC-MS, UV-VIS spectrophotometry (for glycogen, sugar, starch analyses)
- Knowledge of using specialized GPTs (OpenAI Data analyst) and Rstudio in tandem.

References

- Roy, K., Vrba, J., Kajgrova, L. and Mraz, J., 2022. The concept of balanced fish nutrition in temperate European fishponds to tackle eutrophication. *Journal of Cleaner Production*, 364, p.132584.
- Roy, K., Másiľko, J., Kajgrova, L., Kuebutornye, F.K.A., Oberle, M. and Mraz, J., 2023. End-of-season supplementary feeding in European carp ponds with appropriate plant protein and carbohydrate combinations to ecologically boost productivity: Lupine, rapeseed and, triticale. *Aquaculture*, 577, p.739906.
- Roy, K., Kajgrova, L., Capkova, L., Zabransky, L., Petraskova, E., Dvorak, P., Nahlik, V., Kuebutornye, F.K.A., Blabolil, P., Blaha, M. and Vrba, J., 2024. Synergistic digestibility effect by planktonic natural food and habitat renders high digestion efficiency in agastric aquatic consumers. *Science of the Total Environment*, 927, p.172105.
- Roy, K., Vrba, J., Kuebutornye, F.K., Dvorak, P., Kajgrova, L. and Mraz, J., 2024. Fish stocks as phosphorus sources or sinks: Influenced by nutritional and metabolic variations, not solely by dietary content and stoichiometry. *Science of The Total Environment*, 938, p.173611.
- Roy, K., Verdegem, M.C. and Mraz, J., 2025. Ecological Restoration of Inland Aquaculture in Land-Locked Europe: The Role of Semi-Intensive Fishponds and Multitrophic Technologies in Transforming Food Systems. *Reviews in Aquaculture*, 17(1), p.e12999.
- Kuebutornye, F.K.A., Tellbüscher, A.A., Dvořák, P., Roy, K. and Mraz, J., 2025. Feeding value of low opportunity cost biomasses (agri-food by-products) for development of circular pond fish feeds: An evaluation with common carp (*Cyprinus carpio*). *Aquaculture Reports*, 42, p.102753.
- Mraz, J., Mandal, B., Vrba, J., Kajgrova, L., Blaha, M., Kuebutornye, F.K.A., Zabransky, L., Nahlik, V., Lepic, P. and Roy, K., 2025. Nutritional requirement of European fishponds to achieve low and efficient feed use avoiding eutrophication. *Aquaculture*, p.743389.

Funding

Horizon Europe project DIVERSIFY4Future (Project: 101286791); 06/2026-05/2030 (PI: Roy)
INTERREG VA Austria – Czech Republic project: ATCZ00278 – CARMA; 4/2026–12/2028 (PI: Roy)
Other sources: NAZV Kocour QL26010354, OP JAK Mraz CZ.02.01.01/00/23_021/0012616.



Supervisor: Assoc. prof. Vlastimil Stejskal

Contact

E-mail: stejskal@frov.jcu.cz

Tel. +420 389 034 681

Effects of alternative feed ingredients on flesh quality, organoleptic properties, physicochemical characteristics and shelf life of farmed fish

Vliv alternativních krmných surovin na kvalitu svaloviny, senzorické (organoleptické) vlastnosti, fyzikálně-chemické parametry a skladovatelnost ryb

DSP: Rybnářství / Fishery

Annotation

The increasing demand for alternative feed ingredients, including insect meals, fermented plant proteins, plant protein isolates, mycoproteins, and bacterial (single-cell) proteins has stimulated extensive research into their nutritional value and environmental sustainability in aquafeeds (Tran et al., 2022). While considerable attention has been devoted to evaluating growth performance, nutrient digestibility, and immune responses, the effects of these novel ingredients on final product quality remain comparatively underexplored. From a consumer perspective, the acceptability of farmed fish is primarily determined by sensory attributes such as colour, odour, texture, and flavour, alongside physicochemical parameters including pH, water-holding capacity, lipid oxidation, and fatty acid composition. These factors collectively influence shelf life and overall marketability (Alfnès et al., 2006; Gasco et al., 2019; Stejskal et al., 2019; Stejskal et al., 2011; Vácha et al., 2013).

Existing evidence suggests that the inclusion of insect meals can modify fillet fatty acid profiles and alter sensory characteristics, with high substitution levels occasionally leading to the development of off-flavour (Tran et al., 2025). Fermented plant proteins are known to reduce anti-nutritional factors and improve gut health; however, their downstream effects on muscle biochemistry and post-harvest storage stability remain insufficiently characterized. Likewise, mycoproteins and bacterial single-cell proteins have attracted interest due to their balanced amino acid composition and relatively low environmental footprint (Hooft et al., 2025). Emerging developments also include bacterial protein sources capable of producing natural pigments, which may contribute to desirable fillet coloration and enhance product visual quality when incorporated into aquafeeds. Despite this, their influence on key quality parameters, such as lipid oxidation, textural properties, and microbial spoilage during chilled or frozen storage has not been systematically evaluated across aquaculture species (Alfnès et al., 2006).

In the context of increasing regulatory and market pressure to diversify aquafeed formulations without compromising consumer acceptance, there is a clear need for a comprehensive and systematic assessment of how these alternative protein sources affect flesh quality, organoleptic attributes, physicochemical properties, and product shelf life.

The main hypothesis

- Novel feed ingredients including insect meals, fermented plant proteins, plant protein isolates, mycoproteins, and bacterial meals can partially or fully replace conventional fishmeal and soybean meal without adversely affecting the sensory quality of fish flesh.



- The dietary inclusion of these alternative ingredients influences muscle physicochemical properties, including proximate composition, fatty acid profile, pH, water-holding capacity, and textural characteristics.
- Novel feed ingredients modulate lipid oxidative stability and microbial spoilage dynamics during chilled and frozen storage, thereby affecting product shelf life.
- The extent and direction of these effects are dependent on the type of ingredient, its inclusion level, and the processing or fermentation methods applied.
- Bacterial meals containing pigment-producing compounds (e.g., carotenoids) can enhance fillet coloration and improve visual quality of fish products.

Aim(s) of the Ph.D. thesis

- To assess the impact of these alternative ingredients on the organoleptic properties of fish flesh, including colour, odour, texture, and flavour.
- To determine their influence on key physicochemical characteristics of muscle tissue, including proximate composition, fatty acid profile, pH, water-holding capacity, and textural parameters.
- To evaluate the effects of dietary treatments on product shelf life, with particular emphasis on lipid oxidation (e.g., TBARS, peroxide value), microbial spoilage indicators, and sensory changes during chilled and frozen storage.
- To compare quality outcomes across different ingredient types, inclusion levels, and processing or fermentation methods in order to identify optimal feed formulations that preserve product quality.
- To investigate the potential of pigment-producing bacterial meals (e.g., carotenoid-rich biomass) to enhance fillet coloration and improve visual quality of aquaculture products.

Possible approaches to reach the aims / to verify the hypotheses

- A systematic meta-analysis of published studies reporting the effects of novel feed ingredients on fillet quality parameters.
- Controlled feeding trials will be conducted using selected fish species, in which experimental diets containing graded inclusion levels of insect meals, fermented plant proteins, plant protein isolates, mycoproteins, or bacterial meals are compared with conventional fishmeal/soybean meal-based controls. The inclusion levels will be defined based on prior meta-analysis, selecting maximum inclusion thresholds at which no significant reduction in growth performance has been reported, with the aim of rearing animals to market size under practical production conditions
- Sensory evaluation conducted by trained panels employing descriptive analysis methods, including quantitative descriptive analysis (QDA) and temporal dominance of sensations (TDS), to assess colour, odour, texture, and flavour of both raw and cooked products.
- Comprehensive physicochemical characterization of muscle tissue, including proximate composition, fatty acid profiling, pH, water-holding capacity, cooking loss, instrumental texture analysis, and colour measurement using the CIE Lab* system and Salmofan.
- Assessment of lipid oxidation and shelf life through determination of thiobarbituric acid reactive substances (TBARS), peroxide value, free fatty acids during chilled (4°C) and frozen (-20°C) storage.
- Microbiological analyses, including total viable counts and enumeration of specific spoilage organisms, to characterize microbial spoilage dynamics throughout storage
- Statistical analysis using appropriate models and tests, including general linear models (GLM), analysis of variance (ANOVA), non-parametric tests (e.g., Kruskal–Wallis), post hoc comparisons,



principal component analysis (PCA), and correlation analyses to identify relationships between diet composition and quality attributes.

References

- Alfnes, F., Guttormsen, A.G., Steine, G. and Kolstad, K., 2006. Consumers' willingness to pay for the color of salmon: A choice experiment with real economic incentives. *American Journal of Agricultural Economics* 88: 1050-1061.
- Gasco, L., Biasato, I., Dabbou, S., Schiavone, A. and Gai, F., 2019. Animals fed insect-based diets: state-of-the-art on digestibility, performance and product quality. *Animals* 9: 170.
- Stejskal, V., Matousek, J., Podhorec, P., Prokesova, M., Zajic, T., Mraz, J. 2019. The effect of culture system on proximate composition and amino and fatty acid profiles of peeled *Coregonus peled* fillets, *Journal of Aquatic Food Product Technology*, 28:9, 933-943.
- Stejskal, V., Vejsada, P., Cepák, M., Špička, M., Vácha, F., Kouřil, J., Policar, T., 2011. Sensory and textural attributes and fatty acid profiles of fillets of extensively and intensively farmed Eurasian perch (*Perca fluviatilis* L.). *Food Chemistry* 129: 1054-1059
- Tran, H., Van Doan, H., Stejskal, V., 2022. Environmental consequences of using insect meal as an ingredient in aquafeeds: A systematic view. *Reviews in Aquaculture*, 14(1), pp.237-251.
- Tran, H., Nguyen, T., Øverland, M., Gasco, L., Valente, L., & Stejskal, V. (2025). Influence of dietary black soldier fly (*Hermetia illucens*) meal on the quality attributes of aquaculture fish products: a meta-analysis. *Journal of Insects as Food and Feed*, 11(15): 2352-4588.
- Vácha, F., Stejskal, V., Vejsada, P., Kouřil, J., Hlaváč, D., 2013. Texture profile analyses in tench (*Tinca tinca* L., 1758) from extensive and intensive culture. *Acta Veterinaria Brno* 82: 421–425

Funding

Collaboration with companies producing novel feed ingredients, OP JAK, future projects



Fakulta rybnářství
a ochrany vod
Faculty of Fisheries
and Protection
of Waters

Jihočeská univerzita
v Českých Budějovicích
University of South Bohemia
in České Budějovice
Czech Republic

Hydrobiologický ústav BC AV ČR / Institute of Hydrobiology CAS České Budějovice



Supervisor: Milan Říha, Ph.D.

Contact

E-mail: milan.riha@hbu.cas.cz

Tel.: +420 387 775 855

Spatiotemporal dynamics of fish movement: The role of orientation, temperature, and hydrology in habitat selection

Časoprostorová dynamika pohybů ryb: role orientace, teploty a hydrologických poměrů na výběru habitatu

DSP: Rybářství / Fishery

Annotation

River fragmentation caused by dams, weirs, and hydropower plants represents one of the most critical threats to freshwater biodiversity globally (Birnie-Gauvin et al., 2017). While engineering solutions such as fishways and bypass channels are widely implemented to restore longitudinal connectivity, their biological effectiveness is often surprisingly low (Birnie-Gauvin et al., 2017). A major reason for this failure is the discrepancy between hydraulic design and fish behavior. Traditional management focuses on "swimming performance" (can the fish physically swim through?) (Castro-Santos & Haro, 2010), ignoring the critical aspect of "orientation and navigation" (can the fish find the entrance?). Without understanding the mechanisms determining how fish select and utilize their environment (Nathan et al., 2008), conservation efforts in regulated rivers like the Danube remain inefficient.

To improve passage efficiency, we must first understand the baseline spatial strategies of fish in unfragmented or semi-natural environments. Current knowledge of how fish perceive space, maintain headings, and select specific habitats is limited, often relying on coarse-scale data (Hussey et al., 2015). This thesis utilizes high-resolution telemetry from lentic systems (reservoirs) to establish a fundamental understanding of how navigation underpins habitat selection (Rodríguez et al., 2021). By analyzing fine-scale trajectories of model species (e.g., pikeperch *Sander lucioperca*, wels catfish *Silurus glanis*, pike *Esox lucius*), it is possible to define the natural rules of habitat use (how fish search for resting or foraging sites and respond to environmental cues), which serves as a necessary control for understanding their behavior in anthropogenically altered systems (Edelhoff et al., 2016; Říha et al., 2022).

In regulated rivers, these natural selection strategies confront artificial barriers. The complex hydro-morphology of the Danube (Gabčíkovo-Čunovo system) creates a "sensory trap" where altered flow regimes and depth profiles disrupt migration routes (Kubala et al., 2019). Here, the challenge shifts from theoretical orientation to practical passability. Rheophilic fish species must not only possess the motivation to migrate but also the ability to select the correct path towards bypass entrances amidst turbulent tailraces. The problem is exacerbated by the lack of knowledge regarding vertical microhabitat selection; a fish preferring near-bottom structures may simply miss a surface-oriented fishway, regardless of its motivation.

This doctoral research addresses the urgent need to integrate behavioral ecology into river management. By combining the study of fundamental navigation principles with applied telemetry of barrier crossing success, the project aims to identify the specific behavioral bottlenecks limiting connectivity. The analysis of depth selection, thermal habitat preferences, and passage efficiency at key obstacles will provide direct recommendations for optimizing fishway attraction flows. Ultimately, the thesis demonstrates that



successful restoration of migration corridors requires a holistic approach that considers not just the physical permeability of barriers, but the spatiotemporal dynamics of habitat selection of the species attempting to cross them.

The main hypothesis

- In featureless lentic environments, fish rely on allocentric cognitive maps. We hypothesize that navigation towards known habitats is precise and goal-directed regardless of the starting position, significantly exceeding the efficiency of random walk or simple beaconing models.
- In lotic systems, vertical space use is determined by species-specific ecomorphological traits. We hypothesize strict vertical niche partitioning: while pelagic predators as asp (*Leuciscus aspius*) utilize the upper water column, benthic species as starlet (*Acipenser ruthenus*) or barbell (*Barbus barbus*) are obligatorily bound to the riverbed boundary layer, causing them to approach migration barriers at significantly different depth strata. It can caused species-specific differences in barrier permeability.

Aim(s) of the Ph.D. thesis

- To characterize fine-scale movement trajectories and homing precision of different fish species in reservoirs, aiming to confirm the existence of map-like navigation and identify the behavioral rules used for orientation in the absence of flow.
- To assess the spatiotemporal dynamics of rheophilic species (*Acipenser ruthenus*, *Leuciscus aspius*, *Barbus barbus*) in the Danube, with a specific focus on how vertical space use (depth selection) changes in response to discharge, temperature, and migration phase.
- To quantify the passability of the Gabčíkovo-Čunovo waterworks for the target species and identify behavioral bottlenecks (e.g., depth mismatch at fishway entrances) that limit successful migration.
- To quantify the passability of the Gabčíkovo-Čunovo waterworks in relation to species-specific vertical distribution, and to identify behavioral bottlenecks caused by the mismatch between the preferred swimming depths and the vertical position of fishway entrances.

Possible approaches to reach the aims / to verify the hypotheses

- Data Acquisition: Combine high-resolution 3D telemetry (reservoirs) with large-scale linear arrays equipped with depth and temperature sensors (Danube).
- Statistical Framework: Perform all analyses in R (e.g., *actel*, *moveHMM*) using Hidden Markov Models to segment behavioral states, circular statistics for navigation precision, and GLMMs to test abiotic drivers.
- Habitat Modelling: Apply Habitat Selection Models (HSM), specifically Resource Selection Functions (RSF) or Step Selection Functions (SSF), to quantify the selection of microhabitats (depth strata, flow classes) relative to their availability.
- Barrier Assessment: Evaluate passage efficiency using event-time analysis and compare fish vertical profiles with fishway hydraulics to identify migration bottlenecks.



References

- Birnie-Gauvin, K., Aarestrup, K., Riis, T. M., Jepsen, N., & Koed, A. (2017). Shining a light on the loss of rheophilic fish habitat in lowland rivers as a forgotten consequence of barriers, and its implications for management. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 27(6), 1345–1349.
- Castro-Santos, T., & Haro, A. (2010). Fish guidance and passage at barriers. In: *Fish locomotion: an ethological perspective*, 62-89.
- Edelhoff, H., Signer, J., & Balkenhol, N. (2016). Path segmentation for beginners: an overview of current methods for detecting changes in animal movement patterns. *Movement Ecology*, 4(1), 21.
- Hussey, N. E., Kessel, S. T., Aarestrup, K., Cooke, S. J., Cowley, P. D., Fisk, A. T., ... & Whoriskey, F. G. (2015). Aquatic animal telemetry: a panoramic window into the underwater world. *Science*, 348(6240), 1255642.
- Kubala, M., Farský, M., & Pekárik, L. (2019). Migration patterns of sterlet (*Acipenser ruthenus*) in the Middle Danube. *Knowledge & Management of Aquatic Ecosystems*, 420, 40.
- Nathan, R., Getz, W. M., Revilla, E., Holyoak, M., Kadmon, R., Saltz, D., & Smouse, P. E. (2008). A movement ecology paradigm for unifying organismal movement research. *Proceedings of the National Academy of Sciences*, 105(49), 19052–19059.
- Říha, M., Rabaneda-Bueno, R., Jarić, I., Souza, A. T., Vejřík, L., Draštík, V., ... & Peterka, J. (2022). Seasonal habitat use of three predatory fishes in a freshwater ecosystem. *Hydrobiologia*, 849(15), 3351–3371.
- Rodríguez, F., et al. (2021). Spatial cognition in teleost fish: strategies and mechanisms. *Animals*, 11(8), 2271.