



Ph.D. thesis topics 2026/2027

DSP Ochrana vodních ekosystémů / Protection of aquatic ecosystems

Výzkumný ústav rybářský a hydrobiologický / Research Institute of fish Culture and Hydrobiology	
Vodňany	2
Supervisor: Assoc. Prof. Miloš Buřič	3
Responsible compounds in chemical communication of crayfish	3
Supervisor: Ganna Fedorova, Ph.D.	6
Nature-inspired approaches for wastewater treatment and reuse.....	6



Fakulta rybnářství
a ochrany vod
Faculty of Fisheries
and Protection
of Waters

Jihočeská univerzita
v Českých Budějovicích
University of South Bohemia
in České Budějovice
Czech Republic

Výzkumný ústav rybářský a hydrobiologický / Research Institute of fish Culture and Hydrobiology Vodňany





Supervisor: Assoc. Prof. Miloš Buřič

Contact

E-mail: buric@frov.jcu.cz

Tel.: +420 724 204 706

Responsible compounds in chemical communication of crayfish

Sloučeniny odpovědné za chemickou komunikaci raků

DSP: Ochrana vodních ekosystémů / Protection of aquatic ecosystems

Annotation

Recent studies have shown that biogenic amines, such as serotonin, dopamine and octopamine, affect fighting behaviour and dominance status in crayfish. Our own studies have revealed that chemical communication via urine-borne signals is necessary for crayfish to decide whether to fight. The proposed topic should combine known facts about the role of biogenic amines in urinary signals, not only in agonistic interactions or the decision to fight, but also in other behavioural features of crayfish, such as courtship, maternal care and offspring recognition. The same chemical information would act differently in different situations and for different individuals (e.g. juveniles, adults, berrying females, dominant or subordinate). Similarly, different chemical signals would have different priorities in crayfish interactions. Therefore, the same chemical compound can act as an attractant in one case and a repellent in another, or for another individual. Crayfish seem to be a great experimental model species, as they are probably the most easily cultured crustaceans, and their social interactions and frequently used chemical communication make them ideal subjects for study. The results could help to define the role of specific signals in specific behavioural changes. Last but not least, this work could be a further step towards identifying specific compounds (aside from those already known) that are active in crustacean communication, and their potential application in future studies or as attractants in species regulation efforts.

The main hypothesis

- Biogenic amines present in urine-borne chemical signals modulate crayfish social behaviour in a context-dependent manner. The concentration and composition of serotonin, dopamine, and octopamine in chemical signals influence behavioural decisions related to agonistic interactions, courtship, maternal care, and offspring recognition.
- The behavioural effect of the same chemical compound depends on the physiological and social status of the receiving individual. The same urine-borne chemical signal elicits different behavioural responses in juvenile and adult crayfish, dominant and subordinate individuals, or berrying females.
- Chemical signals involved in crayfish communication exhibit hierarchical priority during social interactions. In situations involving competing behavioural motivations (e.g., aggression versus reproduction), some chemical compounds exert stronger influence on behavioural outcomes than others.
- Crayfish chemical communication involves additional behaviourally active compounds beyond currently known biogenic amines. Previously unidentified compounds present in urine-borne signals and significantly contribute to behavioural modulation.



Aim(s) of the Ph.D. thesis

- To characterize the role of biogenic amines in urine-borne chemical communication across different behavioural contexts in crayfish. This objective will assess how serotonin, dopamine, and octopamine are associated with agonistic interactions, courtship, maternal care, and offspring recognition.
- To investigate how physiological condition, developmental stage, and social status influence crayfish responses to chemical signals. This objective will compare behavioural responses among different developmental stages of crayfish exposed to identical chemical cues.
- To determine the relative importance and prioritization of chemical signals during crayfish social interactions. This objective will experimentally evaluate how competing chemical signals influence behavioural decisions e.g. in parental behaviour.
- To confirm the presence (or identify) novel compounds involved in crayfish chemical communication. This objective will focus on detecting previously unknown behaviourally active compounds.

Possible approaches to reach the aims / to verify the hypotheses

- Experimental laboratory work including manipulation and maintenance of crayfish, manipulative experiments in laboratory conditions, pilot field experiments, and injection procedures using specific biogenic amines and their antagonists. Sampling of water, hemolymph and urine for further analysis, elaboration of literature review and analyse the results. Field observations using video-cameras at the places with co-occurrence of invasive and protected/important species.

References

- Bovbjerg, R.V., 1953. Dominance order in the crayfish *Orconectes virilis* (Hagen). *Physiol. Zool.* 26, 173-178.
- Breithaupt, T., Thiel, M., 2010. Chemical communication in crustaceans.
- Breithaupt, T., Eger, P., 2002. Urine makes the difference: chemical communication in fighting crayfish made visible. *J. Exp. Biol.* 205, 1221-1231.
- Breithaupt, T., Gherardi, F., Aquiloni, L., Tricarico, E., 2016. Chemical ecology of crayfish. In: Longshaw M., Stebbing P., *Biology and Ecology of Crayfish*. pp. 140-178.
- Figler, M.H., Twum, M., Finkelstein, J.E., Peeke, H.V.S., Twum, M., 1995. Maternal Aggression in Red Swamp Crayfish (*Procambarus Clarkii*, Girard): The Relation Between Reproductive Status and Outcome of Aggressive Encounters with Male and Female Conspecifics. *Behaviour* 132, 107–125.
- Fossat, P., Bacqué-Cazenave, J., De Deurwaerdère, P., Delbecq, J.P., Cattaert, D., 2014. Anxiety-like behavior in crayfish is controlled by serotonin. *Science* 344, 1293-1297.
- Gherardi, F., Acquistapace, P., Hazlett, B.A., Whisson, G., 2002. Behavioral responses to alarm odors in indigenous and non-indigenous crayfish species: a case study from Western Australia. *Mar. Freshw. Res.* 53, 93-98.
- Gherardi, F., Aquiloni, L., Diéguez-Uribeondo, J., Tricarico, E., 2011. Managing invasive crayfish: is there a hope? *Aquat. Sci.* 73, 185-200.
- Gherardi, F., 2002. Behaviour. In: D.M. Holdich (ed.). *Biology of Freshwater Crayfish*. Blackwell Science, Oxford, pp. 258–290.
- Hazlett, B.A., 2000. Information use by an invading species: do invaders respond more to alarm odors than native species? *Biol. Invasions* 2, 289-294.
- Hazlett, B.A., 1983. Parental behavior in decapod Crustacea. In: *Studies in Adaptation. The Behavior of Higher Crustacea*. pp. 171-193.
- Hazlett, B.A., 2003. Predator recognition and learned irrelevance in the crayfish *Orconectes virilis*. *Ethol.*



109, 765-780.

- Huber, R., Delago, A., 1998. Serotonin alters decisions to withdraw in fighting crayfish, *Astacus astacus*: the motivational concept revisited. *J. Comp. Physiol. A.* 182, 573-583.
- Huber, R., Kravitz, E.A., 1995. A quantitative analysis of agonistic behavior in juvenile American lobsters (*Homarus americanus* L.). *Brain Behav. Evol.* 46, 72-83.
- Huber, R., Smith, K., Delago, A., Isaksson, K., Kravitz, E.A., 1997. Serotonin and aggressive motivation in crustaceans: altering the decision to retreat. *Proc. Natl. Acad. Sci.* 94, 5939-5942.
- Little, E.E., 1975. Chemical communication in maternal behaviour of crayfish. *Nat.* 255, 400-401.
- Little, E.E., 1976. Ontogeny of maternal behavior and brood pheromone in crayfish. *J. Comp. Physiol.* 112, 133-142.
- Mathews, L.M., 2011. Mother—offspring recognition and kin-preferential behaviour in the crayfish *Orconectes limosus*. *Behaviour*, pp. 71-87.
- Stebbing, P.D., Watson, G.J., Fraser, D., Jennings, R., Sibley, P.J., Bentley, M.G., Rushton, S.P., 2004. Evaluation of the capacity of pheromones for the control of invasive non-native crayfish. *English Nature Research Reports*.
- Stebbing, P.D., Watson, G.J., Bentley, M.J., Fraser, D., Jennings, R., Sibley, P.J., 2005. Evaluation of the capacity of pheromones for control of invasive non-native crayfish: part 2. *English Nature Research Reports*.
- Stebbing, P.D., Watson, G.J., Fraser, D., Jennings, R., Sibley, P.J., Bentley, M.G., Rushton, S.P., 2004. Evaluation of the capacity of pheromones for the control of invasive non-native crayfish. *English Nature Research Reports*.
- Stebbing, P.D., Watson, G.J., Bentley, M.J., Fraser, D., Jennings, R., Sibley, P.J., 2005. Evaluation of the capacity of pheromones for control of invasive non-native crayfish: part 2. *English Nature Research Reports*.



Supervisor: Ganna Fedorova, Ph.D.

Contact

E-mail: gfedorova@frov.jcu.cz

Tel.: +420 389 034 756

Nature-inspired approaches for wastewater treatment and reuse

Přírodou inspirované přístupy k čištění a opětovnému použití odpadních vod

DSP: Ochrana vodních ekosystémů / Protection of aquatic ecosystems

Annotation

Every day we are using water and releasing many chemicals to wastewater. With the growing global concern over water scarcity and pollution, there is a critical need for innovative approaches to wastewater treatment that not only effectively remove contaminants but also recover valuable resources from wastewater streams.

Conventional wastewater treatment is not effective enough to remove pollutants from the wastewater, thus wastewater effluents are recognized as the main source of micropollutants in the aquatic environment. Presence of a wide range of emerging contaminants (such as pharmaceuticals, personal care products, perfluorinated compounds) in wastewater treatment plants effluents was reported.

Intensive research is going on to improve elimination of micropollutants from wastewater. So called green approaches, which promote sustainable and environmentally friendly methods for treating wastewater while also reducing the environmental impact of traditional wastewater treatment processes are gaining a lot of attention nowadays. Natural treatment methods include, for example, the use of stabilization ponds for wastewater treatment. This is a cheap and relatively effective way to reduce the content of nutrients and micropollutants in wastewater. Constructed wetlands are also used for wastewater treatment and have the potential to remove a wide range of micropollutants.

Treated wastewater represents a sustainable water source that could be used for fish farming. However, the reuse of treated wastewater poses certain safety risks. Until now, attention has been paid to the risk of microbial contamination or the accumulation of toxic metals in fish produced this way. However, less information is available on the possible contamination of fish with micropollutants.

The main hypothesis

- Additional wastewater treatment steps can improve quality of wastewater effluents
- Water plants can contribute to the elimination of micropollutants from wastewater
- Wastewater reuse for the fish production can be beneficial using the proper practice of fish handling

Aim(s) of the Ph.D. thesis

- To assess the effectiveness of stabilisation pond and constructed wetlands for the elimination of micropollutants from wastewater.
- To evaluate the bioaccumulation of micropollutants in water plants.
- To analyze the content of micropollutants in fish produced in the pond fed with treated wastewater. In case these substances will be detected, to monitor the duration of their purification after transferring the fish into clean water.



Possible approaches to reach the aims / to verify the hypotheses

- Collection of water, plant and fish samples; development and validation of extraction procedures.
- LC/MS analysis of micropollutants in collected samples.
- Data analysis, risk assessment.

References

- Fedorova, G., R. Grabic, K. Grabicova, J. Turek, T. Van Nguyen, T. Randak, B. W. Brooks and V. Zlabek (2022). "Water reuse for aquaculture: Comparative removal efficacy and aquatic hazard reduction of pharmaceuticals by a pond treatment system during a one year study." *Journal of Hazardous Materials* 421.
- Mackuřák, T., M. Mosný, J. řkubák, R. Grabic and L. Birořová (2015). "Fate of psychoactive compounds in wastewater treatment plant and the possibility of their degradation using aquatic plants." *Environ Toxicol Pharmacol* 39(2): 969-973.
- Zaibel, I., D. Zilberg, L. Groisman and S. Arnon (2016). "Impact of treated wastewater reuse and floods on water quality and fish health within a water reservoir in an arid climate." *Science of the Total Environment* 559: 268-281.

CENAKVA Research program

RP2 "New" pollutants in the environment and their effect on freshwater ecosystems